

# EXCLUSIONS PERIODS FOR SPECIFIC DISEASES AND INFECTION

## DIOCESE OF WAGGA WAGGA

Policy Number	11/111
Policy Name	Exclusions Periods
Applicability	CSO and School Personnel
Contact Person	Director of Schools
Policy Status	Approved
Date of Approval	December 2011
Date Last Amended	December 2011
Related Policies/Documents	<ul style="list-style-type: none"> <li>• First Aid and Medication Policy &amp; Procedures</li> <li>• Student Health Care Guidelines (First Aid)</li> </ul>
Review Period:	2013

## RECOMMENDED MINIMUM EXCLUSION PERIODS FOR INFECTIOUS CONDITIONS FOR SCHOOLS, PRE-SCHOOLS AND CHILD CARE CENTRES

National Health and Medical Research Council – December 2005

CONDITION	EXCLUSION OF CASE	EXCLUSION OF CONTACTS
Amoebiasis ( <i>Entamoeba histolytica</i> )	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Campylobacter	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Candidiasis	See 'Thrush'	
Chickenpox (Varicella)	Exclude until all blisters have dried. This is usually at least 5 days after the rash first appeared in unimmunised children and less in immunised children.	Any child with an immune deficiency (for example, leukaemia) or receiving chemotherapy should be excluded for their own protection. Otherwise, not excluded.
CMV (Cytomegalovirus infection)	Exclusion is NOT necessary	Not excluded
Conjunctivitis	Exclude until the discharge from the eyes has stopped unless doctor has diagnosed a non-infectious conjunctivitis	Not excluded
Cryptosporidium infection	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Diarrhoea (No organism identified)	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Diphtheria	Exclude until medical certificate of recovery is received following at least 2 negative throat swabs, the first swab not less than 24 hours after finishing a course of antibiotics followed by another swab 48 hours later.	Exclude contacts that live in the same house until cleared to return by an appropriate health authority.
German measles	See 'Rubella'	
Giardiasis	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Glandular fever (Mononucleosis, EBV infection)	Exclusion is NOT necessary	Not excluded
Hand, foot and mouth disease	Exclude until all blisters have dried.	Not excluded
<i>Haemophilus influenzae</i> type b (Hib)	Exclude until the person has received appropriate antibiotic treatment for at least 4 days.	Not excluded
Head lice (Pediculosis)	Exclusion is NOT necessary if effective treatment is commenced prior to the next day at child care (ie the child doesn't need to be sent home immediately if head lice are detected).	Not excluded
Hepatitis A	Exclude until a medical certificate of recovery is received, but not before seven days after the onset of jaundice.	Not excluded
Hepatitis B	Exclusion is NOT necessary	Not excluded
Hepatitis C	Exclusion is NOT necessary	Not excluded
Herpes simplex (cold sores, fever blisters)	Exclusion is not necessary if the person is developmentally capable of maintaining hygiene practices to minimise the risk of transmission. If the person is unable to comply with these practices they should be excluded until the sores are dry. Sores should be covered by a dressing where possible.	Not excluded
Human Immunodeficiency Virus (HIV/AIDS)	Exclusion is NOT necessary. If the person is severely immunocompromised, they will be vulnerable to other people's illnesses.	Not excluded

<b>CONDITION</b>	<b>EXCLUSION OF CASE</b>	<b>EXCLUSION OF CONTACTS</b>
Hydatid disease	Exclusion is NOT necessary	Not excluded
Impetigo (school sores)	Exclude until appropriate antibiotic treatment has commenced. Any sores on exposed skin should be covered with a watertight dressing.	Not excluded
Influenza and influenza like illnesses	Exclude until well	Not excluded
Legionnaires' disease	Exclusion is NOT necessary	Not excluded
Leprosy	Exclude until approval to return has been given by an appropriate health authority	Not excluded
Measles	Exclude for 4 days after the onset of the rash	Immunised and immune contacts are not excluded. Non-immunised contacts of a case are to be excluded from child care until 14 days after the first day of appearance of rash in the last case, unless immunised within 72 hours of first contact during the infectious period with the first case. All immunocompromised children should be excluded until 14 days after the first day of appearance of rash in the last case.
Meningitis (bacterial)	Exclude until well and has received appropriate antibiotics	Not excluded
Meningitis (viral)	Exclude until well	Not excluded
Meningococcal infection	Exclude until appropriate antibiotic treatment has been completed	Not excluded
Molluscum contagiosum	Exclusion is NOT necessary	Not excluded
Mumps	Exclude for nine days after onset of swelling	Not excluded
Norovirus	Exclude until there has not been a loose bowel motion or vomiting for 48 hours	Not excluded
Parvovirus infection (fifth disease, erythema infectiosum, slapped cheek syndrome)	Exclusion is NOT necessary	Not excluded
Pertussis	See 'Whooping Cough'	
Respiratory Syncytial virus	Exclusion is NOT necessary	Not excluded
Ringworm/tinea	Exclude until the day after appropriate antifungal treatment has commenced	Not excluded
Roseola	Exclusion is NOT necessary	Not excluded
Ross River virus	Exclusion is NOT necessary	Not excluded
Rotavirus infection	Children are to be excluded from the centre until there has not been a loose bowel motion or vomiting for 24 hours	Not excluded
Rubella (German measles)	Exclude until fully recovered or for at least four days after the onset of the rash	Not excluded
Salmonella infection	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Scabies	Exclude until the day after appropriate treatment has commenced	Not excluded
Scarlet fever	See 'Streptococcal sore throat'	
School sores	See 'Impetigo'	
Shigella infection	Exclude until there has not been a loose bowel motion for 24 hours	Not excluded
Streptococcal sore throat (including scarlet fever)	Exclude until the person has received antibiotic treatment for at least 24 hours and feels well	Not excluded
Thrush (candidiasis)	Exclusion is NOT necessary	Not excluded
Toxoplasmosis	Exclusion is NOT necessary	Not excluded

<b>CONDITION</b>	<b>EXCLUSION OF CASE</b>	<b>EXCLUSION OF CONTACTS</b>
Tuberculosis (TB)	Exclude until medical certificate is produced from an appropriate health authority	Not excluded
Typhoid, Paratyphoid	Exclude until medical certificate is produced from appropriate health authority	Not excluded unless considered necessary by public health authorities
Varicella	See 'Chickenpox'	
Viral gastroenteritis (viral diarrhoea)	Children are to be excluded from the centre until there has not been a loose bowel motion or vomiting for 24 hours	Not excluded
Warts	Exclusion is NOT necessary	Not excluded
Whooping cough (pertussis)	Exclude until five days after starting appropriate antibiotic treatment or for 21 days from the onset of coughing	Contacts that live in the same house as the case and have received less than three doses of pertussis vaccine are to be excluded from the centre until they have had 5 days of an appropriate course of antibiotics. If antibiotics have not been taken, these contacts must be excluded for 21 days after their last exposure to the case while the person was infectious.
Worms	Exclusion not necessary if treatment has occurred	Not excluded

### **Staying Healthy in Child Care – Preventing infectious diseases in child care 4<sup>th</sup> edition**